

**INDICATIONS:** For topical use on animals as an aid in reducing pain, licking and chewing on wounds and surgical incisions. Kills up to 99% of surface germs in 15 seconds or less that can potentially cause an infection.

**ACTIVE INGREDIENTS**

Povidone-Iodine (2% available iodine)  
Bitrex™ Denatonium Benzoate  
Lidocaine 2% w/v  
Ethyl Alcohol – less than 80%  
Isopropyl Alcohol – less than 5%

**PURPOSE**

Antiseptic  
Bitter Agent  
Pain Relief  
Antiseptic  
Antiseptic

**DIRECTIONS:** Pour into a spray bottle dialed to spray pattern, hold sprayer about 4–6 inches from area to be treated. Spray 1 or more times — allow to dry. Repeat as needed. Forms a highly visible coating that is durable.

**WARNINGS:** Do not use in the eyes. Do not use if allergic or sensitive to the active ingredients. Prolonged exposure may cause irritation. If swallowed, get medical help or contact a Poison Control Center right away. Avoid inhalation.

Store upright at 15°–25° C (59°–77° F). Brief excursions up to 40° C (104° F) are permitted.

NDC 51072-036-01

**INDICACIONES:** Para uso tópico en animales para ayudar a reducir el dolor, lametazo, y masticación en heridas e incisiones quirúrgicas. Mata hasta el 99% de los gérmenes superficiales en 15 segundos o menos, que potencialmente puede causar una infección.

**INGREDIENTES ACTIVOS**

Yodopovidona (Yodo al 2%)  
Bitrex™ Benzoato de denatonio  
Lidocaina 2% p/v  
Alcohol etílico – menos del 80%  
Alcohol isopropílico – menos del 5%

**PROPÓSITO**

Antiséptico  
Agente amargo  
Anestésico local  
Antiséptico  
Antiséptico

**INSTRUCCIONES DE USO:**

Verter en botella de spray, marcado a patrón de rociado, mantener pulverizador a unas 4–6 pulgadas de la zona a tratar. Rociar una o más veces, dejar a secar. Repetir según sea necesario. Forma una capa que es altamente visible y duradera.

**ADVERTENCIAS:** No utilice en los ojos. No utilice en caso de alergia o de sensibilidad a los ingredientes activos. La exposición prolongada puede causar irritación. En caso de ingestión, buscar ayuda médica o contactar al centro de Control de Venenos (Poison Control Center) inmediatamente. Evitar la inhalación.

Almacenar verticalmente a 15°–25° C (59°–77° F). Excursiones breves hasta 40° C (104° F) son admisibles.

# BARRIER®

## Wound Care Spray With Pain Relief 2% Available Iodine With Lidocaine

*(Patent pending)*

For Animal Use Only

### 1 Gallon (3.79 liters)



MANUFACTURED IN THE USA



**FLAMMABLE — USE WITH CAUTION  
FOR EXTERNAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
WET HAIR OR FUR IS FLAMMABLE  
DOT Name: Ethanol Solutions UN1170,3,PGII**

REORDER NO: 19002

MANUFACTURED BY:  
Aurora Pharmaceutical, Inc.  
NORTHFIELD,  
MINNESOTA 55057  
1-888-215-1256  
aurorapharmaceutical.com  
IN 50-1083 06/2023



3 51072 03601 6

TAKE TIME



OBSERVE LABEL  
DIRECTIONS