

INDICATIONS FOR USE: CoccidAid® (amprolium oral solution) 9.6% Oral Solution is intended for the treatment of coccidiosis in growing chickens, turkeys, and laying hens. If no improvement is noted within 3 days, have the diagnosis confirmed and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.

USE DIRECTIONS: Give amprolium at the 0.012% level (8 fl oz per 50 gallons) as soon as coccidiosis is diagnosed and continue for 3 to 5 days. (In severe outbreaks, give amprolium at the 0.024% level.) Continue with 0.006% amprolium medicated water for an additional 1 to 2 weeks. No other source of drinking water should be available to the birds during this time. Use this as the sole source of amprolium. To disperse particles in the product, shake the bottle prior to each use.

▶ **WITHDRAWAL PERIODS:** No withdrawal period is required when used according to labeling. ◀

USER SAFETY WARNING: Keep this and all drugs out of the reach of children. NOT FOR HUMAN USE.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY.

MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

Restricted Drug (California) - Use Only as Directed

STORAGE: Store between 5°-25°C (41°- 77°F) with brief excursions to 40°C. Benzoic acid 0.1% added as preservative.

CONTACT INFORMATION: To report suspected adverse drug experiences, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Aurora Pharmaceutical at 1-888-215-1256 or www.aurorapharmaceutical.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae



REORDER NO: 24001

Aurora Pharmaceutical, Inc.
NORTHFIELD, MN 55057
1-888-215-1256

www.aurorapharmaceutical.com

IN 50-1465 09/2024

TAKE TIME



OBSERVE LABEL
DIRECTIONS



NDC 51072-090-00



CocciAid® (amprolium oral solution) 9.6% Oral Solution Coccidiostat

**Water-soluble treatment for coccidiosis
Treats 800 gallons at 0.012% level**

Approved by FDA under ANADA # 200-630

Net Contents:

1 Gallon (128 fl. oz) (3.785 L)

Active Ingredient:

amprolium.....9.6%



FOR ANIMAL USE ONLY

To Prepare 50 Gallons of Medicated Water

DOSAGE LEVEL	MIXING DIRECTIONS
0.024%	Add 1 pint (16 fluid ounces) of CocciAid® (amprolium oral solution) 9.6% Oral Solution to about 5 gallons of water in a 50-gallon medication barrel. Stir, then add water to the 50-gallon mark. <i>Stir thoroughly.</i>
0.012%	Follow same directions as above but use 1/2 pint (8 fluid ounces) of CocciAid® 9.6% Oral Solution.
0.006%	Follow same directions as above but use 4 fluid ounces of CocciAid® 9.6% Oral Solution.

For Automatic Water Proportioners

For automatic water proportioners that meter 1 fluid ounce of stock solution per gallon of drinking water.

DOSAGE LEVEL	COCCIAID® 9.6% ORAL SOLUTION PER GALLON OF STOCK SOLUTION
0.024%	41 fl oz
0.012%	20.5 fl oz
0.006%	10.25 fl oz

Note: Make drinking water fresh daily. Stock solutions for proportioners may be stored in a clean, closed labeled container for up to 3 days.

CocciAid® is a registered trademark of Aurora Pharmaceutical, Inc.