

**ACTIVE INGREDIENT**

Sodium Salicylate .....48.6% w/v

NDC 51072-038-01

**CALF LABEL CLAIM**

**Indications For Use, Calves:** Supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (e.g., anti-infective) therapy if necessary.

**Contraindications, Calves:** Do not use Sodium Salicylate in neonates or calves less than 2 weeks of age.

**Dosage, Calves:** 40 mg Sodium Salicylate per kg body weight (4 mL per 100 lbs.) once daily, for 1–3 days. Administer orally in drinking water or milk (replacer).

**SWINE AND POULTRY LABEL CLAIM**

**Indications:** For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

*(continued on next panel)*

REORDER NO: 21004

MANUFACTURED BY:

**Aurora Pharmaceutical, Inc.**  
NORTHFIELD, MINNESOTA 55057  
1-888-215-1256  
www.aurorapharmaceutical.com  
IN 50-1108 12/2024



TAKE TIME



OBSERVE LABEL DIRECTIONS



MANUFACTURED IN THE USA

**ORAL-PRO®**

# Sodium Salicylate Concentrate 48.6% w/v

For Use in Livestock Only

**Keep Out of Reach of Children**

Net Contents:

**1 Gallon (3.79 Liters)****DIRECTIONS FOR USE****For Analgesic and Antipyretic Use**

**Water Proportioner Use:** Add 8 ounces (236 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

**Livestock Tank Use:** Add 8 ounces (236 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

**For Anti-Inflammatory/Anti-Prostaglandin Use — Day 1**

**Water Proportioner Use:** Add 16 ounces (473 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

**Livestock Tank Use:** Add 16 ounces (473 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

**Day 2 through 7**

**Water Proportioner Use:** Add 10 ounces (296 mL) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

**Livestock Tank Use:** Add 10 ounces (296 mL) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Prepare fresh solutions daily. Repeat as necessary.

**WARNING**

Not for use in lactating dairy cattle.

Do not administer concentrated solution by direct oral administration — gastrointestinal irritation or overdose may occur. Do not use in piglets less than 3 weeks of age.

**STORAGE**

Store upright up to 25°C (77°F) with brief excursions to 40°C (104°F).

Do not use if allergic or sensitive to the active ingredients.

**CAUTION**

Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product.

Gradual darkening will not affect product stability.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.