

ACTIVE INGREDIENTS

Sodium Salicylate..... 60% w/v
 Caffeine5.7% w/v

NDC 51072-039-01

DIRECTIONS FOR USE *(continued)***For Anti-Inflammatory/Anti-Prostaglandin Use — Day 1**

Water Proportioner Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Day 2 through 7

Water Proportioner Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Livestock Tank Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily. Prepare fresh solutions daily. Repeat as necessary.

WARNING

Not for use in lactating dairy cattle.

Do not administer concentrated solution by direct oral administration — gastro-intestinal irritation or overdose may occur. Do not use in piglets less than 3 weeks of age.

Store upright up to 25°C (77°F) with brief excursions to 40°C (104°F).

Do not use if allergic or sensitive to the active ingredients.

CAUTION

Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution, or place in warm water. Gently invert container to ensure uniformity of product.

Gradual darkening will not affect product stability.

ORAL-PRO®

Sodium Salicylate Concentrate 60% w/v with Caffeine 5.7% w/v

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:

1 Gallon (3.79 Liters)

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

DIRECTIONS FOR USE:**For Analgesic and Antipyretic Use**

Water Proportioner Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate with Caffeine 5.7% to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

*(continued on next panel)***MANUFACTURED BY:**

Aurora Pharmaceutical, Inc.
 NORTHFIELD, MINNESOTA 55057
 1-888-215-1256
www.aurorapharmaceutical.com
 IN 50-1110 12/2024

REORDER NO: 21003

TAKE TIME



OBSERVE LABEL DIRECTIONS

**MANUFACTURED
 IN THE USA****Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.